

Adverse childhood experiences from family and society contribute to increased risk of depressive symptoms and cognitive impairment: a cross-sectional study

Ziyang Ren ^{1,2}, Yanan Luo,³ Xiaoying Zheng,⁴ Jufen Liu ^{1,2}

To cite: Ren Z, Luo Y, Zheng X, *et al.* Adverse childhood experiences from family and society contribute to increased risk of depressive symptoms and cognitive impairment: a cross-sectional study. *General Psychiatry* 2023;**36**:e101039. doi:10.1136/gpsych-2023-101039

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/gpsych-2023-101039>).

Received 16 February 2023
Accepted 15 August 2023



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Jufen Liu;
liujufen@bjmu.edu.cn

Xiaoying Zheng;
zhengxiaoying@sph.pumc.edu.cn

ABSTRACT

Background Family environments can shape children's personalities and social networks, rendering distinguishing adverse childhood experiences (ACEs) from family and society essential, but related evidence remains limited.

Aims This cross-sectional study aimed to investigate the correlations between intrafamilial and social ACEs, their associations with depressive symptoms and cognitive impairment and the (education-moderated) mediating role of social ACEs.

Methods Data for this cross-sectional study were from the China Health and Retirement Longitudinal Study. Nine intrafamilial (0, 1, 2, 3, and 4 or more) and three social (0, 1, and 2 or more) ACEs were identified. Depressive symptoms were assessed using the 10-item Center for Epidemiological Studies Depression Scale. Global cognition, including episodic memory and mental intactness, was calculated as z scores. Binary and ordered logistic regressions, generalised linear models with Gaussian family and identity link, and mediation analysis were used.

Results 13 435 participants aged 59.0 (51.0–66.0) were included. Compared with participants with no intrafamilial ACEs, those with 1, 2, 3, and 4 or more intrafamilial ACEs tended to develop more social ACEs, with odds ratios (ORs) of 1.55 (95% confidence interval (CI): 1.36 to 1.76), 2.36 (95% CI: 2.08 to 2.68), 3.46 (95% CI: 3.02 to 3.96) and 6.10 (95% CI: 5.30 to 7.02), respectively. Both intrafamilial and social ACEs were associated with depressive symptoms (OR >3 for four or more intrafamilial ACEs and two or more social ACEs) and global cognition ($\beta = -0.26$ for four or more intrafamilial ACEs and $\beta = -0.29$ for two or more social ACEs). Social ACEs mediated the associations of intrafamilial ACEs with depressive symptoms and global cognition by 12.3% and 13.1%, respectively. Furthermore, as education levels increased, the impact of intrafamilial ACEs on depressive symptoms was increasingly mediated through social ACEs, while the mediating role of social ACEs between intrafamilial ACEs and cognitive impairment gradually diminished.

Conclusions Improving children's social environments and elevating general education can prevent later-life depressive symptoms and cognitive impairment attributed to ACEs in China.

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Adverse childhood experiences (ACEs), generally classified as deprivation and threat, are known to induce depression and cognitive impairment in later life. According to the ecological systems theory, family environments can influence children's personalities and social networks. Thus, taking a sociological perspective when studying ACEs, for instance, the mediating role of social ACEs among the associations of intrafamilial ACEs with depressive symptoms and cognitive impairment, seems worthy of investigation. Furthermore, exploring whether education modifies the mediating role of social ACEs can provide population evidence for reducing the health burdens attributed to ACEs, but related evidence remains controversial.

WHAT THIS STUDY ADDS

⇒ This study found significant correlations between intrafamilial and social ACEs, as well as their significant associations with depressive symptoms and cognitive impairment. Of all ACEs, parental mental illness, loneliness and unfriendly neighbours exhibited the highest depressive and cognitive risks. Social ACEs mediated more than 10% of the associations of intrafamilial ACEs with depressive symptoms and cognitive impairment. Interestingly, as education levels increased, the impact of intrafamilial ACEs on depressive symptoms and cognitive impairment gradually weakened or even diminished.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ Given the high prevalence of ACEs in China, our findings highlight that parents, teachers and community workers should concentrate on preventing children's occurrence of social ACEs and fostering the development of their social abilities while working to reduce ACEs within the family. In addition, the Chinese government should strive to raise the educational levels for all to prevent widespread ACE-induced depressive symptoms and cognitive impairment.

INTRODUCTION

Due to the rapid expansion of the ageing population, mental and memory-related disorders, such as depression and dementia, presently affect 13.3% and 7.0% of the elderly population worldwide and have escalated into significant public health issues.^{1,2} Therefore, identifying potential lifelong risk factors for these disorders is essential to alleviate their considerable socioeconomic and medical burdens.

Adverse childhood experiences (ACEs) encompass a broad range of traumatic events children and adolescents may encounter, precipitating subsequent risks of depression and cognitive impairment in their adult lives.³ Given their distinct effects on neurodevelopment, ACEs are typically categorised as deprivation and threat.⁴ Nevertheless, considering the impact of intrafamilial circumstances on the development of children's personalities and social relationships, a sociological perspective becomes indispensable when studying ACEs.^{5,6} It helps to comprehensively understand ACEs and facilitates the formulation of efficacious prevention and intervention strategies, particularly in China, where such evidence remains limited.

A previous study has shown that peer bullying during childhood is a mediator between intrafamilial aggression and depression in adulthood, suggesting that children growing up in unfavourable familial environments may experience social adversities in their early years, leading to long-term health issues.⁷ However, this study only focused on maltreatment as intrafamilial adversity and peer bullying as social adversity. The associations between broader intrafamilial ACEs, such as emotional neglect and economic hardship, and social ACEs, such as loneliness and neighbourhood environments, have not been extensively investigated thus far. Even though individuals with ACEs are susceptible to developing depression and cognitive impairment,³ evidence is limited concerning the distinctions between intrafamilial and social ACEs and whether social ACEs mediate the impact of intrafamilial ACEs on health issues. Furthermore, research conducted in the UK has provided evidence suggesting a negative association between ACEs and education. Years of education are a significant component of socioeconomic status and cognitive reserve known to benefit mental health and cognitive function.^{8,9} However, a recent study conducted in China has reported no significant moderated effects of education among the association of ACEs with depression and cognitive impairment,¹⁰ leading to controversial findings. Given the gap in older adults' education and the substantial cultural differences between China and Western countries, investigating how education moderates the mediating role of social ACEs between intrafamilial ACEs and health outcomes may help develop targeted interventions to reduce the health hazards of ACEs in older Chinese.

To fill in these research gaps, we used the China Health and Retirement Longitudinal Study (CHARLS) to investigate the associations of intrafamilial and social ACEs with depressive symptoms and cognitive impairment,

the mediating role of social ACEs and the education-moderated mediating role of social ACEs in middle-aged and older Chinese.

METHODS

Study population

For this cross-sectional study, we used data from CHARLS, a nationally representative survey that included adults aged 45 years and older from 450 villages and urban communities across 28 provinces in China. The detailed procedures taken by CHARLS have been described elsewhere.¹¹ After the 2011 baseline survey, there were follow-ups in 2013, 2015 and 2018. In 2014, CHARLS additionally carried out a life history survey to retrospectively record the respondents' life experiences from birth. CHARLS 2014 and 2015 waves were used in this study.

Among 21 095 participants recruited in 2015, those who were aged <45 or with incomplete data on age, gender, residence, education, marital status, smoking history, drinking history and history of chronic diseases ($n=1787$) and who had incomplete data on ACEs ($n=5873$) were excluded, leaving 13 435 participants included at baseline (figure 1).

Definition of ACEs

The detailed definitions of ACEs are shown in online supplemental table S1. In CHARLS, ACEs before the age of 17 were assessed using the 2014 Life History Questionnaire. Nine intrafamilial ACEs (emotional neglect, family violence, parental separation or divorce, parental behavioural problem, parental mental illness, parental disability, parental death, physical abuse and economic adversity) and three social ACEs (bullying, loneliness and unfriendly neighbours) were finally identified. All ACEs were dichotomised and summed to obtain intrafamilial ACEs and social ACEs, with values ranging from 0 to 9 and 0 to 3, respectively. Intrafamilial ACEs were then divided into 0, 1, 2, 3, and 4 or more; social ACEs were classified as 0, 1, and 2 or more.

Assessment of depressive symptoms

CHARLS uses the 10-item Center for Epidemiological Studies Depression Scale to measure depressive symptoms.¹² Participants were asked about their feelings for 10 aspects during the last week, such as feeling bothered, having trouble concentrating and feeling depressed, with options including (1) rarely or none of the time (1 day); (2) some or a little of the time (1–2 days); (3) sometimes or a significant amount of the time (3–4 days); and (4) most or all of the time (5–7 days). These four choices were then valued as 0, 1, 2 and 3 in sequence, with a total depressive score ranging from 0 to 30. A higher depressive score indicates more depressive symptoms. Individuals with a depressive score of ≥ 10 were defined as having depressive symptoms.¹³

Assessment of cognitive function

The cognitive function includes episodic memory and mental intactness (orientation, attention and




Figure 1 Flowchart and design of the study. ACEs, adverse childhood experiences. CHARLS, China Health and Retirement Longitudinal Study.

visuoconstruction) in CHARLS, with a total cognitive score ranging from 0 to 21.¹⁰ A higher cognitive score indicates better cognitive function. The average of immediate (within 2 min) and delayed (4–10 min later) word recalls using 10 random words was adopted to measure episodic memory (0–10). Orientation was defined by asking about the day, week, month, season and year of the interview (0–5). Attention was determined using five serial subtractions of 7 from 100 (0–5). Visuoconstruction was defined by asking participants to redraw a previously displayed figure (0–1). Global cognition, episodic memory and mental intactness were then calculated as each-5-year-age stratified z scores. All cognitive z scores were further normalised using Blom transformation to fit Gaussian distributions.

Covariates

Information on age, gender, residence, education, marital status, smoking history, drinking history and number of diseases was collected through face-to-face interviews. Residence was classified as rural and urban. Education was divided into primary school or less, middle school, and high school or above. Marital status was classified as married/cohabiting or single. Smoking and drinking history were classified as never or ever. Number of diseases was defined as using self-reported physician diagnoses or using treatments and was divided into 0, 1, and 2 or more.

Statistical analysis

The baseline characteristics of included participants were described as medians and interquartile ranges (IQRs) for continuous variables, given their skewed

distribution, and frequency and per cent (%) for categorical variables. To compare characteristics across participants with different numbers of intrafamilial ACEs, differences were assessed by Kruskal-Wallis tests (continuous or ordered categorical variables) and χ^2 tests (binary variables).

Binary logistic regression was used to investigate the association of each and overall intrafamilial ACEs with each social ACE. Ordered logistic regression was further used to investigate the association of each and overall intrafamilial ACEs with overall social ACEs. All models were adjusted for age, gender, residence, smoking history, drinking history and number of diseases.

Furthermore, the fully adjusted associations of intrafamilial and social ACEs with depressive symptoms were investigated using binary logistic regression, while the associations with cognitive function (global cognition, episodic memory and mental intactness) were investigated using generalised linear models with Gaussian family and identity link. Corresponding associations of each ACE with depressive symptoms and cognitive function were investigated using similar models. Gender, residence and marital status-stratified analyses were further conducted.

Before mediation analysis, the correlation between intrafamilial ACEs and social ACEs was examined using Spearman's rank correlation, and the variance inflation factors (VIFs) were also calculated to assess potential collinearity issues. The Spearman's rank correlation was 0.27 (<0.40), and the VIFs for intrafamilial ACEs and social ACEs were 1.10 and 1.09, respectively. Also, the

Table 1 Baseline characteristics of included participants (n=13 435)

Characteristics	Number of intrafamilial ACEs				χ^2	P value	
	0 (n=2745)	1 (n=3938)	2 (n=3197)	3 (n=1997)			4 or more (n=1558)
Age, year	58.0 (51.0–66.0)	59.0 (51.0–66.0)	59.0 (52.0–66.0)	60.0 (52.0–66.0)	59.0 (52.0–66.0)	9.63	0.047
Gender						42.26	<0.001
Men	1191 (43.4)	1893 (48.1)	1637 (51.2)	1007 (50.4)	775 (49.7)		
Women	1554 (56.6)	2045 (51.9)	1560 (48.8)	990 (49.6)	783 (50.3)		
Residence						37.67	<0.001
Rural	1604 (58.4)	2310 (58.7)	2012 (62.9)	1250 (62.6)	1022 (65.6)		
Urban	1141 (41.6)	1628 (41.3)	1185 (37.1)	747 (37.4)	536 (34.4)		
Education						132.29	<0.001
Primary school or less	1726 (62.9)	2609 (66.3)	2209 (69.1)	1460 (73.1)	1208 (77.5)		
Middle school	651 (23.7)	819 (20.8)	671 (21.0)	344 (17.2)	240 (15.4)		
High school or above	368 (13.4)	510 (13.0)	317 (9.9)	193 (9.7)	110 (7.1)		
Marital status						9.46	0.051
Married/cohabiting	2446 (89.1)	3505 (89.0)	2821 (88.2)	1733 (86.8)	1361 (87.4)		
Single	299 (10.9)	433 (11.0)	376 (11.8)	264 (13.2)	197 (12.6)		
Social ACEs						908.22	<0.001
0	2309 (84.1)	3045 (77.3)	2204 (68.9)	1208 (60.5)	739 (47.4)		
1	376 (13.7)	730 (18.5)	779 (24.4)	580 (29.0)	535 (34.3)		
2 or more	60 (2.2)	163 (4.1)	214 (6.7)	209 (10.5)	284 (18.2)		
Smoking history						41.35	<0.001
Ever	1073 (39.1)	1731 (44.0)	1486 (46.5)	932 (46.7)	703 (45.1)		
Never	1672 (60.9)	2207 (56.0)	1711 (53.5)	1065 (53.3)	855 (54.9)		
Drinking history						29.74	<0.001
Ever	890 (32.4)	1402 (35.6)	1220 (38.2)	762 (38.2)	600 (38.5)		
Never	1855 (67.6)	2536 (64.4)	1977 (61.8)	1235 (61.8)	958 (61.5)		
Number of chronic diseases						150.20	<0.001
0	937 (34.1)	1247 (31.7)	908 (28.4)	520 (26.0)	365 (23.4)		
1	800 (29.1)	1121 (28.5)	888 (27.8)	533 (26.7)	349 (22.4)		
2 or more	1008 (36.7)	1570 (39.9)	1401 (43.8)	944 (47.3)	844 (54.2)		
Depressive symptoms*						435.94	<0.001
Yes	578 (22.8)	948 (26.3)	930 (32.2)	745 (41.4)	697 (50.0)		
No	1958 (77.2)	2661 (73.7)	1961 (67.8)	1053 (58.6)	696 (50.0)		

Continued

Table 1 Continued

Characteristics	Number of intrafamilial ACEs					χ^2	P value
	0 (n=2745)	1 (n=3938)	2 (n=3197)	3 (n=1997)	4 or more (n=1558)		
Global cognition*	12.5 (9.5–15.0)	12.0 (9.5–14.5)	12.0 (9.0–14.5)	11.5 (9.0–14.5)	11.5 (8.0–14.0)	60.66	<0.001
Episodic memory*	4.0 (2.5–5.0)	4.0 (2.5–5.0)	3.5 (2.5–5.0)	3.5 (2.5–5.0)	3.5 (2.0–4.5)	36.06	<0.001
Mental intactness*	9.0 (6.0–11.0)	8.0 (6.0–10.0)	8.0 (6.0–10.0)	8.0 (6.0–10.0)	7.0 (6.0–10.0)	51.09	<0.001

Values are presented as n (%) or median (IQR).
 *The analytical sample is less than 13 435.
 ACEs, adverse childhood experiences; IQR, interquartile range.

VIFs of all other variables were less than 10, suggesting no severe collinearity.

Subsequently, the mediating role of social ACEs in the associations of intrafamilial ACEs with depressive symptoms and cognitive function (global cognition, episodic memory and mental intactness), as well as the education-moderated mediating role of social ACEs in these associations, was explored based on the ‘mediation’ package in R. For instance, the equations for exploring the mediating role of social ACEs between intrafamilial ACEs and depressive symptoms were as follows:

Model 1: Social ACEs=Intrafamilial ACEs+Covariates (ordered logistic regression)

Model 2: Depressive symptoms=Intrafamilial ACEs+Social ACEs+Covariates (binary logistic regression)

Reporting of this study was done under the Strengthening the Reporting of Observational Studies in Epidemiology guidelines. Analyses were performed using R statistical software V.4.2.3 (R Project for Statistical Computing). All analyses were two-sided, and a p value of <0.05, a 95% confidence interval (CI) of odds ratio (OR) that did not cross 1.00 or a 95% CI of β that did not cross 0 was considered statistically significant.

RESULTS

The baseline characteristics of the included participants are shown in table 1. A total of 6503 (48.4%) men and 6932 (51.6%) women aged 59.0 (51.0–66.0) were included. Those with more intrafamilial ACEs tend to develop more social ACEs, depressive symptoms and cognitive impairment (all p values <0.001).

According to table 2, when compared with those with no intrafamilial ACEs, participants with more overall intrafamilial ACEs tend to develop more overall social ACEs, with ORs of 1.55 (95% CI: 1.36 to 1.76), 2.36 (95% CI: 2.08 to 2.68), 3.46 (95% CI: 3.02 to 3.96) and 6.10 (95% CI: 5.30 to 7.02) for 1, 2, 3, and 4 or more intrafamilial ACEs, respectively.

Both intrafamilial and social ACEs are associated with depressive symptoms, with ORs of 3.23 (95% CI: 2.79 to 3.73) for four or more intrafamilial ACEs and 3.15 (95% CI: 2.71 to 3.67) for two or more social ACEs (online supplemental table S2) when compared with those with no corresponding ACEs. Similarly, both intrafamilial and social ACEs are negatively associated with global cognition (β =−0.26, 95% CI: −0.33 to −0.19 for four or more intrafamilial ACEs and β =−0.29, 95% CI: −0.36 to −0.21 for two or more social ACEs), episodic memory (β =−0.16, 95% CI: −0.23 to −0.09 for four or more intrafamilial ACEs and β =−0.19, 95% CI: −0.27 to −0.11 for two or more social ACEs) and mental intactness (β =−0.26, 95% CI: −0.33 to −0.19 for four or more intrafamilial ACEs and β =−0.27, 95% CI: −0.35 to −0.19 for two or more social ACEs). Online supplemental table S3 suggests that parental mental illness, loneliness and unfriendly neighbours are associated the most with depressive symptoms and cognitive impairment among all ACEs. Furthermore,

Table 2 Associations between intrafamilial ACEs and social ACEs: binary and ordered logistic regressions

Exposure	Outcome			
	Bullying	Loneliness	Unfriendly neighbours	Overall social ACEs
	OR (95% CI)			
Specific intrafamilial ACEs				
Emotional neglect	1.11 (1.01 to 1.23)	1.05 (0.94 to 1.18)	1.22 (1.09 to 1.37)	1.13 (1.04 to 1.22)
Family violence	2.42 (2.20 to 2.68)	1.57 (1.40 to 1.76)	1.40 (1.25 to 1.58)	1.94 (1.79 to 2.11)
Parental separation or divorce	2.46 (1.48 to 4.07)	1.82 (1.01 to 3.29)	1.38 (0.72 to 2.63)	1.81 (1.15 to 2.85)
Parental behavioural problem	1.54 (1.25 to 1.91)	1.69 (1.34 to 2.13)	1.28 (0.99 to 1.65)	1.62 (1.36 to 1.93)
Parental mental illness	2.27 (2.00 to 2.57)	2.91 (2.55 to 3.32)	2.32 (2.02 to 2.65)	2.74 (2.48 to 3.04)
Parental disability	1.59 (1.43 to 1.77)	1.75 (1.55 to 1.97)	1.50 (1.32 to 1.69)	1.70 (1.56 to 1.85)
Parental death	1.12 (0.97 to 1.30)	1.85 (1.61 to 2.14)	1.44 (1.24 to 1.68)	1.45 (1.30 to 1.62)
Physical abuse	2.82 (2.55 to 3.10)	1.56 (1.39 to 1.75)	1.43 (1.27 to 1.60)	2.08 (1.92 to 2.25)
Economic adversity	2.09 (1.90 to 2.30)	1.96 (1.76 to 2.19)	2.43 (2.17 to 2.71)	2.28 (2.12 to 2.46)
Overall intrafamilial ACEs				
0	1.00 (Reference)	1.00 (Reference)	1.00 (Reference)	1.00 (Reference)
1	1.65 (1.37 to 1.98)	1.50 (1.23 to 1.83)	1.43 (1.18 to 1.74)	1.55 (1.36 to 1.76)
2	2.80 (2.34 to 3.35)	1.92 (1.58 to 2.34)	2.00 (1.65 to 2.43)	2.36 (2.08 to 2.68)
3	4.27 (3.55 to 5.14)	2.71 (2.21 to 3.32)	2.60 (2.12 to 3.19)	3.46 (3.02 to 3.96)
4 or more	6.50 (5.39 to 7.83)	4.80 (3.93 to 5.86)	4.06 (3.32 to 4.97)	6.10 (5.30 to 7.02)

Models were adjusted for age, gender, residence, marital status, smoking history, drinking history and number of chronic diseases. ACEs, adverse childhood experiences; CI, confidence interval; OR, odds ratio.

the associations of intrafamilial and social ACEs with depressive symptoms and cognitive impairment remain significant in sex, residence and marital status-stratified analysis, as shown in online supplemental tables S4–S6.

As shown in [figure 2](#), social ACEs mediate the associations of intrafamilial ACEs with depressive symptoms, global cognition, episodic memory and mental intactness by 12.3%, 13.1%, 9.8% and 15.5%, respectively. However, the mediating role of social ACEs between intrafamilial ACEs and episodic memory was not so robust in stratified analysis. According to the final moderated mediation models in [figure 3](#), significant interactions of intrafamilial ACEs and education towards depressive symptoms and of social ACEs and education towards cognitive impairment were found. Online supplemental table S7 provides additional evidence showing a noteworthy decrease in the direct effects of intrafamilial ACEs on depressive symptoms while an increase in the mediation proportion of social ACEs as the level of education increases. Furthermore, the direct and indirect effects of intrafamilial ACEs and the mediation proportion of social ACEs towards global cognition, episodic memory and mental intactness become non-significant in those with an education level of high school or above (online supplemental table S7).

DISCUSSION

Main findings

In this retrospective study, we found that intrafamilial ACEs may lead to social ACEs and, thus, further facilitate

depressive symptoms and cognitive impairment in adulthood. Interestingly, as education levels increased, the impact of intrafamilial ACEs on depressive symptoms was increasingly mediated through social ACEs; however, the mediating role of social ACEs between intrafamilial ACEs and cognitive impairment gradually diminished and became non-significant among older adults with high school or above education levels.

Prior research has consistently demonstrated that ACEs are associated with increased risks of later-life mental disorders.^{3,14} Increased chronic inflammation and dysfunctional hypothalamic-pituitary-adrenal axis can be used to interpret the associations between ACEs and depression.¹⁵ As for cognitive impairment, the blockage of neurodevelopment by ACEs may play a role.^{16,17} Importantly, we further confirmed that social ACEs mediated the associations of intrafamilial ACEs with both depressive symptoms and cognitive impairment. Family environments have been found to influence children's behaviour and personality development.¹⁸ Previous research has shown that children who grow up in chaotic homes are more likely to cheat in games.¹⁹ Harsh parenting, quite prevalent in China, can also lead to child aggression.²⁰ Some theories can be used to interpret our findings. According to the ecological systems theory, the family represents the most fundamental microsystem in children's lives and plays a significant role in their development and adaptation.²¹ Intrafamilial ACEs, such as family conflict or parental relationship breakdown, can disrupt




Figure 2 The mediating role of social ACEs in the associations of intrafamilial ACEs with depressive symptoms and cognitive function. Models were adjusted for age, gender, residence, marital status, smoking history, drinking history and number of chronic diseases. ACEs, adverse childhood experiences.

children’s microsystems, leading to adverse consequences for their socialisation process, such as feelings of insecurity and unacceptance. Simultaneously, social learning theory emphasises that children acquire behaviours by observing and imitating the adults and peers around them.²² Hence, intrafamilial ACEs can influence children’s ability to develop healthy relationships with peers. For instance, conflict and violence within the family may shape children’s perception of violence as an effective means of problem-solving, increasing their likelihood of engaging in aggressive behaviour and encountering conflict within their social environment.

Among all ACEs, parental mental illness, loneliness and unfriendly neighbours are most strongly associated with depressive symptoms and cognitive impairment. According to our findings, parental mental illness may contribute to other intrafamilial ACEs like physical abuse²² and is associated with social ACEs the most among all

intrafamilial ACEs. In addition, to some extent, parental mental illness may indicate a genetic predisposition for children’s mental illness.²³ Loneliness during childhood is a relatively direct reflection of children’s mental health. Furthermore, living in unfriendly neighbourhoods shows substantial associations with depressive symptoms and cognitive impairment later in life. In unfriendly neighbourhoods, children may experience feelings of isolation and have limited access to emotional support, which can contribute to developing depressive symptoms.²⁴ Unfriendly neighbourhoods may also provide inadequate opportunities for cognitive stimulation, intellectual engagement or positive social interactions, resulting in children’s weaker cognitive foundations and reduced confidence in social interactions.²⁵

Interestingly, the adverse impacts of intrafamilial ACEs on depressive symptoms and cognitive impairment gradually diminish with elevated education. Higher levels




Figure 3 The final moderated mediation models. Models were adjusted for age, gender, residence, marital status, smoking history, drinking history and number of chronic diseases. * $p < 0.05$. ACEs, adverse childhood experiences.

of education generally provide individuals with more resources that can help mitigate the effects of ACEs. For instance, highly educated individuals often have better access to healthcare, mental health services and social support networks,²⁶ all of which can offer crucial support in coping with the emotional and cognitive challenges attributed to ACEs. Higher education levels are also associated with greater resilience and autonomy.²⁷ Education equips individuals with knowledge, skills and self-confidence, enabling them to navigate the difficulties posed by ACEs effectively. Additionally, higher education provides opportunities for cognitive stimulation and intellectual growth. These mental skills act as protective factors against the detrimental effects of ACEs on depressive symptoms and cognitive impairment.²⁸

In this retrospective study, we comprehensively explored the associations of intrafamilial and social ACEs in China. We indicated for the first time the mediating role of social ACEs in the associations of intrafamilial ACEs with depressive symptoms and cognitive impairment. Our findings highlight the necessity of distinguishing different ACE types via a more sociological perspective. Furthermore, the education-moderated mediation analyses indicate that elevating education levels can reduce ACEs' adverse effects on depressive symptoms and cognitive impairment. The national representativeness and the rigorous procedures of CHARLS ensure the robustness of our conclusions.

Limitations

Our study has several limitations. First, the ACEs data were self-reported, which could lead to recall bias. However, this strategy has been used consistently in most previous studies and seems difficult to avoid. In addition, the study design was retrospective rather than cohort, leaving the causal association less established. Finally, there are racial differences among the effects of ACEs, according to Assari²⁹ and Assari and Lankarani.³⁰

However, information on ethnicity in CHARLS is unavailable, rendering its effect modification in this study not explored.

Implications

In addition to resolving conflicts within their families, our study emphasises that interventions for children with intrafamilial ACEs should concentrate on preventing the occurrence of social ACEs and fostering the development of their social abilities. This should be done with the combined efforts of parents, teachers and community workers. Moreover, China should strive to elevate general education levels to prevent and reduce depressive symptoms and cognitive impairment attributed to prevalent ACEs.

Author affiliations

¹Department of Epidemiology and Biostatistics, School of Public Health, Peking University, Beijing, China

²Institute of Reproductive and Child Health/National Health Commission Key Laboratory of Reproductive Health, Peking University, Beijing, China

³Department of Global Health, School of Public Health, Peking University, Beijing, China

⁴School of Population Medicine and Public Health, Chinese Academy of Medical Sciences/Peking Union Medical College, Beijing, China

Contributors JL and ZR designed the study. ZR managed and analysed the data. ZR prepared the first draft. ZR reviewed and edited the manuscript, with comments from YL, XZ and JL. All authors were involved in revising the paper. JL accepted full responsibility for the work and the conduct of the study, had access to the data, and controlled the decision to publish.

Funding This study was funded by the Major Project of the National Social Science Fund of China (21ZDA107) and the Fundamental Research Funds for the Central Universities (7101303357).

Disclaimer The funding source had no involvement in the study planning, execution or reporting.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants and ethical approval for all CHARLS waves was granted from the Institutional Review Board at Peking University. The IRB approval number for the main household survey, including

anthropometrics, is IRB00001052-11015. The IRB approval number for biomarker collection is IRB00001052-11014. All participants provided written informed consent. Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Ziyang Ren <http://orcid.org/0000-0003-0223-4073>

Jufen Liu <http://orcid.org/0000-0001-5844-8973>

REFERENCES

- Abdoli N, Salari N, Darvishi N, *et al*. The global prevalence of major depressive disorder (MDD) among the elderly: a systematic review and meta-analysis. *Neurosci Biobehav Rev* 2022;132:1067–73.
- Cao Q, Tan C-C, Xu W, *et al*. The prevalence of dementia: a systematic review and meta-analysis. *J Alzheimers Dis* 2020;73:1157–66.
- Nelson CA, Scott RD, Bhutta ZA, *et al*. Adversity in childhood is linked to mental and physical health throughout life. *BMJ* 2020;371:m3048.
- McLaughlin KA, Sheridan MA, Lambert HK. Childhood adversity and neural development: deprivation and threat as distinct dimensions of early experience. *Neurosci Biobehav Rev* 2014;47:578–91.
- Haslam Z, Taylor EP. The relationship between child neglect and adolescent interpersonal functioning: a systematic review. *Child Abuse Negl* 2022;125:105510.
- Han W, Chen BB. An evolutionary life history approach to understanding mental health. *Gen Psychiatr* 2020;33:e100113.
- Wang Q. Association of childhood intrafamilial aggression and childhood peer bullying with adult depressive symptoms in China. *JAMA Netw Open* 2020;3:e2012557.
- Houtepen LC, Heron J, Suderman MJ, *et al*. Associations of adverse childhood experiences with educational attainment and adolescent health and the role of family and socioeconomic factors: a prospective cohort study in the UK. *PLoS Med* 2020;17:e1003031.
- Wagg E, Blyth FM, Cumming RG, *et al*. Socioeconomic position and healthy ageing: a systematic review of cross-sectional and longitudinal studies. *Ageing Res Rev* 2021;69:101365.
- Zhang T, Kan L, Jin C, *et al*. Adverse childhood experiences and their impacts on subsequent depression and cognitive impairment in Chinese adults: a nationwide multi-center study. *Journal of Affective Disorders* 2023;323:884–92.
- Zhao Y, Hu Y, Smith JP, *et al*. Cohort profile: the China health and retirement longitudinal study (CHARLS). *Int J Epidemiol* 2014;43:61–8.
- Chen H, Mui AC. Factorial validity of the Center for Epidemiologic Studies Depression Scale short form in older population in China. *Int Psychogeriatr* 2014;26:49–57.
- Li H, Liu X, Zheng Q, *et al*. Gender differences and determinants of late-life depression in China: a cross-sectional study based on CHARLS. *Journal of Affective Disorders* 2022;309:178–85.
- Gu W, Zhao Q, Yuan C, *et al*. Impact of adverse childhood experiences on the symptom severity of different mental disorders: a cross-diagnostic study. *Gen Psychiatr* 2022;35:e100741.
- Yang JZ, Kang CY, Yuan J, *et al*. Effect of adverse childhood experiences on hypothalamic-pituitary-adrenal (HPA) axis function and antidepressant efficacy in untreated first episode patients with major depressive disorder. *Psychoneuroendocrinology* 2021;134:S0306-4530(21)00306-1.
- Sheridan MA, McLaughlin KA. Dimensions of early experience and neural development: deprivation and threat. *Trends Cogn Sci* 2014;18:580–5.
- Lund JI, Boles K, Radford A, *et al*. A systematic review of childhood adversity and executive functions outcomes among adults. *Arch Clin Neuropsychol* 2022;37:1118–32.
- Lucas-Thompson RG, Goldberg WA. Family relationships and children's stress responses. *Adv Child Dev Behav* 2011;40:243–99.
- Vrijhof CI, van der Voort A, van IJendoorn MH, *et al*. Stressful family environments and children's behavioral control: a multimethod test and replication study with twins. *J Fam Psychol* 2018;32:49–59.
- Qi W. Harsh parenting and child aggression: child moral disengagement as the mediator and negative parental attribution as the moderator. *Child Abuse Negl* 2019;91:12–22.
- Kazdin AE. Encyclopedia of psychology, vol. 3. In: *Ecological systems theory*. Washington: Oxford University Press, 2000.
- Bandura A, Walters RH. *Social learning theory*. Englewood cliffs Prentice Hall, 1977.
- Cantwell DP, Baker L. Parental mental illness and psychiatric disorders in "at risk" children. *J Clin Psychiatry* 1984;45:503–7.
- Chen H, Xiong P, Chen L, *et al*. Childhood neighborhood quality, friendship, and risk of depressive symptoms in adults: the China health and retirement longitudinal study. *J Affect Disord* 2020;276:732–7.
- Peng C, Han SH, Burr JA. Perceptions of childhood neighborhood social cohesion and cognitive function in middle and late adulthood. *Gerontologist* 2022;62:1266–77.
- Steele LS, Dewa CS, Lin E, *et al*. Education level, income level and mental health services use in Canada: associations and policy implications. *Healthc Policy* 2007;3:96–106.
- Armfield JM, Ey L-A, Zufferey C, *et al*. Educational strengths and functional resilience at the start of primary school following child maltreatment. *Child Abuse Negl* 2021;122:105301.
- Lövdén M, Fratiglioni L, Glymour MM, *et al*. Education and cognitive functioning across the life span. *Psychol Sci Public Interest* 2020;21:6–41.
- Assari S. Family socioeconomic status and exposure to childhood trauma: racial differences. *Children (Basel)* 2020;7:57.
- Assari S, Lankarani MM. Stressful life events and risk of depression 25 years later: race and gender differences. *Front Public Health* 2016;4:49.



Ziyang Ren graduated from the School of Public Health at Zhejiang University in Zhejiang, China, in June 2022, with a Bachelor of Medicine. He was admitted to the Department of Epidemiology and Biostatistics, School of Public Health, Peking University in China as a PhD student in September 2022. He has authored more than ten research papers and has published as the first author in *JAMA Network Open*, *Journal of Affective Disorders* and other journals. His main research interests include neuropsychiatric disorders, social epidemiology, and life course epidemiology.