Cognitive-behavioural therapy for personal recovery of patients with schizophrenia: A systematic review and meta-analysis

Weiliang Wang,¹ Yuqiu Zhou,¹ Nannan Chai,² Dongwei Liu¹

To cite: Wang W, Zhou Y, Chai N, et al. Cognitive-behavioural therapy for personal recovery of patients with schizophrenia: A systematic review and meta-analysis. *General Psychiatry* 2019;**32**:e100040. doi:10.1136/gpsych-2018-100040

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/gpsych-2018-100040>).

Received 23 November 2018

Revised 03 July 2019

Accepted 17 July 2019

ABSTRACT

Background To date, cognitive-behavioural therapy (CBT) trials have primarily focused on clinical recovery; however, personal recovery is actually the fundamental aspect of the recovery process. The aim of this study was to summarise and synthesise the existing evidence regarding the effectiveness of CBT for personal recovery in patients with schizophrenia.

Aim This study aimed to determine the effectiveness of CBT for personal recovery in patients with schizophrenia.

Methods A systematic search of the literature in PsycINFO, PubMed, Cochrane (CENTRAL), Embase and Web of Science (SCI) was conducted to identify randomised controlled trials reporting the impact of CBT interventions on personal recovery in patients with schizophrenia. The estimated effect sizes of the main study outcomes were calculated to estimate the magnitude of the treatment effects of CBT on personal recovery. We also evaluated the CBT's effect size at the end-of-treatment and long-term (follow-up) changes in time as well as personal recovery.

Results Twenty-five studies were included in the analysis. The effect of CBT on personal recovery was 2.27 (95% CI 0.10 to 2.45; $I^2=0\%$; $p=0.04$) at post-treatment and the long-term effect size was 2.62 (95% CI 0.51 to 4.41; $I^2=0\%$; $p=0.02$). During the post-treatment period, the pooled effect size of CBT was 0.01 (95% CI -0.12 to 0.15; $I^2=33.0\%$; $p>0.05$) for quality of life (QoL), 0.61 (95% CI 0.35 to 1.130; $I^2=30.8\%$; $p<0.01$) for psychosocial health-related QoL, -1.77 (95% CI -3.29 to -0.25; $I^2=40\%$; $p=0.02$) for hopelessness and 1.85 (95% CI 0.69 to 3.01; $I^2=41\%$; $p<0.01$) for self-esteem. We also summarised the effects of CBT on QoL (subscale scores not included in the evaluation of the pooled effect size), self-confidence and connectedness, and all results corresponded to positive effects. However, there was insufficient evidence regarding the long-term effects of CBT on personal recovery.

Conclusions CBT is an effective therapy with meaningful clinical effect sizes on personal recovery and some aspects of personal recovery of schizophrenia after treatment. However, the effect is relatively immediate and rapidly decreases as time progresses. Therefore, in the future, more studies should focus on the mechanism of CBT for personal recovery and the factors that influence the long-term effects of CBT.

Trial registration number CRD42018085643.

INTRODUCTION

Schizophrenia, a severe mental illness, affects more than 21 million people worldwide.¹ The persistent negative symptoms and cognitive impairment associated with schizophrenia have led to its classification among the top 25 leading causes of disability worldwide and the top 11 leading causes of reduced years lived with disability in 2013.² The WHO's Mental Health Action Plan 2013–2020 highlights the steps required to provide appropriate services for people with schizophrenia.³

Cognitive-behavioural therapy (CBT) is the primarily recommended psychological treatment for schizophrenia according to major guidelines.^{4,5} Abundant studies have proven that the effects of CBT on reducing positive symptoms,⁶ improving negative symptoms,⁷ conferring functional improvement,⁸ reducing the time of relapse⁹ and reducing suicidal ideation¹⁰ in patients with schizophrenia are significant. However, the remission of clinical symptoms does not meet the criterion for rehabilitation, and patient organisations have emphasised that recovery can occur even when psychotic symptoms are persistent.¹¹ Recovery is an ongoing, complex and multidimensional process. According to different perspectives, schizophrenia recovery can be classified as clinical and personal.¹² In the treatment of patients with schizophrenia, the primary goal traditionally is the clinical recovery. Clinical recovery includes remission of symptoms and functional improvement, which is the premise of other non-pharmacological treatments and rehabilitations. The relationship between the clinical and personal recovery is somewhat correlated, and both should be considered when monitoring the treatments and outcomes of patients with schizophrenia.¹³ The term 'personal recovery', which based on the perspective of individuals who have

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹School of Nursing, Daqing Campus of Harbin Medical University, Daqing, Heilongjiang, China

²School of Nursing, Chifeng University, Chifeng, Inner Mongolia Autonomous Region, China

Correspondence to

Professor Yuqiu Zhou, Harbin, China; hmuhyx@163.com

experienced mental illness,¹⁴ has been widely used in the literature to describe the patient-based definition of recovery.¹⁵ The most frequently cited patient-based definition is ‘the development of new meaning and purpose in one’s life as one grows beyond the catastrophic effects of mental illness.’¹⁶

Personal recovery varies from person to person, and it is difficult to define common characteristics. Different researchers also have their own definitions of personal recovery. Andresen *et al*¹⁷ concluded that personal recovery included four key points: finding hope; re-establishing identity; finding meaning in life; and taking responsibility for recovery. Leamy *et al*¹⁸ posited that the categories of personal recovery encompass connectedness, hope, identity, meaning and empowerment. Based on a cluster analysis of self-reported personal recovery-related variables, Rossi *et al*¹⁹ identified resilience, self-esteem, coping strategies, stigma and personal strength. Furthermore, quality of life (QoL),^{20–21} taking control of one’s life,^{22–23} personal confidence and reliance on others²⁴ have been found to be important components of personal recovery.

According to a review of the abundant literature on personal recovery, which have most consistently identified connectedness, hope and empowerment as relevant categories,¹³ and discussions by our research team, we decided to use the CHIME personal recovery model defined by Leamy *et al*¹⁸: Connectedness–Hope–Identity–Meaning–Empowerment. In addition, this model is consistent with the context of recovery defined by the WHO.²⁵ Based on the progress of pharmacological treatment of acute psychiatric symptoms of schizophrenia, QoL measurement has become an important indicator for evaluating clinical outcomes in patients with schizophrenia.²⁶ Furthermore, QoL is one of the most commonly used outcome assessments. To better address the concept of personal recovery, we have also considered QoL in the assessment of personal recovery.

The concept of recovery is multidimensional; why, then, should personal recovery be the primary focus? Personal recovery pertains to patient’s ability to live a favourable, dignified and meaningful life. It is the core element of recovery from the patient perspective, and it is the ultimate aim of mental illness treatment. Moreover, some components of personal recovery, such as hope, are the foundations and preconditions of treatment and other outcomes.²⁷ In addition, Jahn *et al*’s²⁸ finding suggests that personal recovery is a protective factor against suicidal ideation in individuals with schizophrenia. However, existing knowledge about the role of CBT in personal recovery is highly limited, and the current meta-analyses focused only on clinical outcomes;^{29–30} therefore, the aim of this study was to determine the effectiveness of CBT for personal recovery in patients with schizophrenia. This research has been registered at PROSPERO (CRD: 42018085643), and the study protocol can be obtained via the following website: <https://www.crd.york.ac.uk/PROSPERO/#recordDetails>.

METHODS

The Preferred Reporting Items for Systematic Reviews and Meta-Analyses statement³¹ were followed in all steps of this research.

Search strategy

Five electronic databases, PsycINFO, PubMed, the Cochrane Library (CENTRAL), Embase and Web of Science (SCI), were searched for relevant papers published before 31 December 2018 with the following search terms: (‘psychosis’ OR ‘psychotic’ OR ‘schizophrenia’ OR ‘schizoaffective disorder’) AND (‘cognitive therapy’ OR ‘cognitive behavior* therapy’ OR ‘cognitive behaviour* therapy’ OR ‘CBT’ AND (‘random’ OR ‘randomized control trial’ OR ‘clinical trial’ OR ‘trial’). Manual searches were also performed by reviewing the reference lists of related papers. Two reviewers (WW and NC) independently screened the search results by reviewing titles and abstracts. The full texts of relevant articles screened in previous step were downloaded for further screening. If the same data were reported in more than one publication, only the paper with the more complete data was included. Any disagreement was settled by discussion with the third author (ZY). (See the flow chart of the study, figure 1.)

Inclusion and exclusion criteria

The definition of personal recovery in our study included five components: connectedness, hope, identity, meaning, empowerment; and QoL. Studies that fulfilled the following criteria were included: (1) randomised controlled trials; (2) publications with full texts written in English; (3) participants diagnosed with schizophrenia or schizophrenia spectrum disorder based on the International Classification of Diseases-Tenth Revision or Diagnostic and Statistical Manual of Mental Disorders-Fourth/Fifth Edition; (4) the use of a valid measure to assess personal recovery (CHIME and QoL); and (5) a psychological intervention of CBT or a CBT-modified programme, but not in combination with other psychological interventions.

The exclusion criteria were as follows: (1) no relevant data available for further analysis; (2) article types other than randomised controlled trials (RCT), such as comments, letters and reviews; and (3) other cognitive therapies, such as cognitive training, cognitive-behavioural social skills training, cognitive remediation therapy or cognitive enhancement therapy.

Data extraction

Data extraction was performed by two independent reviewers (WW and NC) who used a specific worksheet designed before the literature search to minimise errors in data extraction. Data extraction was conducted using the full-text versions of the RCTs. The data regarding basic characteristics and outcome measures, including study identity (first author, publication year and country); study design (randomisation, concealment of allocation,

Figure 1 The flowchart of search and study selection. CBT, cognitive-behavioural therapy; RCT, randomised controlled trial.

blinding); patient characteristics (number of study participants, mean age); intervention characteristics (treatment protocol, length of treatment, number of sessions and type of comparisons); and all relevant outcomes (types of outcome measures, instruments and follow-up periods) were extracted from all included studies.

Quality assessment

The quality of the RCTs enrolled in our study was assessed using the Cochrane Collaboration's tool for determining the risk of bias in randomised trials.³² According to the Cochrane assessment tool, the relevant information was extracted from each study, and the study was rated as 'high risk', 'low risk' or 'unclear risk'. Disagreements were resolved by consensus.

Statistical analysis

All the pooled effect size was performed by RevMan V.5.0. The I^2 statistic was used to evaluate the heterogeneity of the studies. Sensitivity analysis and subgroup analysis were performed to deal with heterogeneity. Forest plots were also drawn to visualise the extent of heterogeneity across studies. Publication bias was evaluated using Egger's test by Stata (V.14.2). Hedge's g was used to determine the effect size of continuous outcomes. Considering the heterogeneity of the personal recovery outcome measures, we summarised the pooled results narratively with descriptive statistics and textual descriptions. A two-tailed $p < 0.05$ was considered statistically significant. A power analysis to examine the reliability of the pooled result was performed with GPower V.3.1.

The Grading of Recommendations Assessment, Development and Evaluation (GRADE) approach was performed to assess confidence of evidence (CoE) for each comparison. CoE of outcomes was rated based on study design, risk of bias, inconsistency of results, indirectness of evidence, imprecision and publication bias. We used the GRADEpro Guideline Development Tool to assess the CoE of the critical outcomes.

RESULTS

Study selection

A total of 4146 articles were retrieved from the electronic databases. After duplicates were discarded, 2493 remaining studies were screened. According to the inclusion and exclusion criteria, approximately 2337 records were removed after screening to determine whether the article titles and abstracts were relevant to the topic of the review. The full texts of the remaining 156 studies were reviewed. We manually searched by reviewing the reference lists from the full-text articles, and none of the references were related to the topic. Finally, 131 studies did not meet the inclusion criteria, leaving 25 studies for inclusion in this review. The details of the search process are shown in [figure 1](#).

Study characteristics

All the participants included in the 25 studies^{33–57} were diagnosed with schizophrenia or schizoaffective disorders; the patients in seven studies were recruited from hospitals, and the patients in the other 18 studies were recruited from the community or mental health centres. The intervention treatment provided in most of the studies was CBT; 1 of the 25 studies⁴⁴ used cognitive-behavioural oriented services (CBOS) as the intervention group's treatment, but CBOS is still considered as CBT on the basis of its core theory and implementation. All the comparison groups received treatment as usual, supportive group, standard care, standard treatment (ST), standard support, befriending, wait-listing or parent psychoeducation therapy, except in two studies^{37,56} that compared CBT versus CBT plus clozapine and CBT versus CBT plus thioridazine. Three studies^{53,54,57} reported directly the personal recovery using the Questionnaire about the Process of Recovery (QPR) scale, 11 studies^{33,35,37,40,43,50,51,53–56} reported QoL; 5 studies^{34,36,45–47} reported hope as an outcome of personal recovery; self-esteem, a core element of a better and more meaningful life for psychiatry patients, was measured in 11 studies^{34,39,41,42,45–49,52,57}; 3 studies^{39,44,55} reported relationships with others as an outcome; and 1 study³⁹ reported self-confidence as an outcome of personal recovery. No studies reported empowerment as an outcome. Twelve of the studies reported sufficient follow-up data to evaluate the long-term effect of CBT in schizophrenia ([table 1](#)).

Quality of the studies

The risk of bias for each study is available in the online supplementary material 1. Eleven^{35,36,38,39,42,48,50,51,53–55} of the 25 studies were universally assessed as having a low risk of bias across all domains. Fifteen trials^{33,36,38,39,42,44,45,48,50,51,53–57} employed adequate methods of sequence generation, 10 trials^{31,33,34,37,40,43,46,47,49,52} were not clear. In addition, the risk of bias due to inadequate allocation concealment was unclear in seven trials,^{34,37,40,43,45,46,52} and four^{33,41,47,49} trials did not include allocation concealment. Lack of blinding of the assessors led to a high risk of bias for some outcomes in four studies,^{41,42,44,47} and an unclear risk in four studies.^{33,37,43,52} A high risk of bias due to lack of participants or staff blinding was found in two studies^{33,47} and was unclear in three studies.^{37,43,52} There was a high risk of bias due to incomplete outcome data for two of the included trials,^{43,45} one trial⁴⁴ did not report all outcomes.

Main efficacy meta-analysis

Primary outcome
Three studies^{53,54,57} reported the effect of CBT for personal recovery measured by QPR. The random effects meta-analysis yielded a summary effect size of 2.27 (95% CI 0.16 to 4.45; $I^2=0\%$; $p=0.04$, power=0.61). Egger's test indicated that there was no publication bias ($p=0.96$, 95% CI -18.1 to 17.96). The long-term effect of CBT was measured in these three studies and the pooled effect size was 2.89 (95% CI 0.51 to 4.47; $I^2=0\%$; $p=0.02$, power=0.13) ([figure 2](#)).

Secondary outcomes

Effect size of QoL

Nine studies^{33,37,40,43,51,53–56} reported QoL total scores based on questionnaires. The random effects meta-analysis yielded a summary effect size of 0.01 (95% CI -0.12 to 0.15; $I^2=33.0\%$; $p>0.05$) and a power analysis result of 0.97 ([figure 2](#)). Egger's test indicated that there was no publication bias ($p=0.54$, 95% CI -1.40 to 2.48). The participants in three studies^{40,43,54} were recruited from hospitals, and the participants in six studies were recruited from outside the hospital. Both the inpatient and outpatient subgroups yielded a small and non-significant effect of schizophrenia on QoL (online supplementary material 1). Seven studies^{33,37,51,53–56} reported CBT follow-up for QoL in schizophrenia, and the pooled effect size was 0.06 (95% CI -0.03 to 0.15; $I^2=15\%$; $p>0.05$) with a small power of 0.19. The follow-up times differed among the studies (1 month;³³ 3 months;^{37,55,56} 6 months and above^{51,53,54}). We evaluated the effect sizes using Cohen's d , and the pooled effect sizes were 0.36 (1 month), 0.08 (95% CI -0.31 to 0.47; $I^2=57.0\%$; $p>0.05$, power=0.87) (3 months) and 0.04 (95% CI -0.00 to 0.09; $I^2=0\%$; $p=0.05$, power=0.11) (6 months and above) respectively. After performing the sensitivity analyses, no substantial change in the new pooled effect size was observed.

Four studies^{35,38–40} reported the psychosocial well-being of the patients, as measured by the Warwick-Edinburgh Mental Well-Being Scale (WEMWBS), a subscale of the

Table 1 Characteristics of the included studies

Study (year)	Place	Mean age, years (I/C)	Inpatient or outpatient	Size (n): (I/C)	Study conditions	Form and dose of intervention*	Personal recovery	Instrument	Follow-up† (month)
Barretto et al, ³³ 2009	Brazil	39.8/33.2	Outpatient	21 12/9	CBT versus BF	Individual; 20 sessions (first 15 sessions weekly, last 5 sessions every other week); 45 min per session	Quality of life	QoL	1
Barrowclough et al, ³⁴ 2006	UK	39/33	Outpatient	113 57/56	CBT versus TAU	Group; 18 sessions over 6 months; 2 hours per session	Hopelessness; self-esteem	BHS; RSE	6
Bechdolf et al, ³⁵ 2010	Germany	32.2/31.4	Inpatient	88 40/48	CBT versus PE	Group; 16 sessions in 8 weeks; 1–1.5 hours per session	Quality of life	MSQoL	4
Birchwood et al, ³⁶ 2014	UK	37.4 38.8/35.9	Outpatient	19 8/9	CBT versus TAU	Group; 25 sessions over 9 months; unclear	Hopelessness	BHS	9
Edwards et al, ³⁷ 2011	Australia	22.0/22.5 20.8/20.5	Outpatient	48 12/11 11/14	CBT versus TDZ CBT versus CLZ	Group; twice weekly for 12 weeks, at least 15 sessions (16 sessions; 19 sessions); unclear	Quality of life	QLS	3
Freeman et al, ³⁸ 2014	UK	41.9/41.5	Outpatient	15 15/15	CBT versus SC	Individual; 6 sessions over 8 weeks; unclear	Self-confidence; well-being; relationships; self-esteem.	BCSS; WEMWBS; SCS; RSQ	1
Freeman et al, ³⁹ 2015	UK	40.9/42.1	Outpatient	150 73/77	CBT versus SC	Individual; 6 sessions over 8 weeks; 1 hour per session	Well-being	WEMWBS	4
van der Gaag et al, ⁴⁰ 2011	Netherlands	36.5/37.4	Inpatient	216 110/106	CBT versus TAU	Group; 26 sessions weekly; unclear	Quality of life	WHO-QoL	–
Garety et al, ⁴¹ 1994	UK	39.6/37.6	Inpatient	20 13/7	CBT versus WL	Individual; 12 sessions weekly over 6 months; unclear	Self-esteem	RSE	–
Gumley et al, ⁴² 2006	UK	35.8/36.7	Outpatient	144 72/72	CBT versus TAU	Group; sessions between entry and 12 weeks; 2 sessions per week delivered at the appearance of early signs of relapse; unclear	Self-esteem	RSE	–
Halperin et al, ⁴³ 2000	Australia	NR	Inpatient	16 7/9	CBT versus WL	Group; 8 sessions over 8 weeks, delivered weekly; 2 hours per session	Quality of life	Q-LES-Q	–
Klingberg et al, ⁴⁴ 2010	Germany	33/33	Inpatient	169 84/85	CBOS versus TAU	Group; 40 weekly sessions weekly, 1 hour per session; 4 fortnightly sessions, 2 hours per session; 6 weekly and 8 fortnightly sessions	Relationships	–	–
Kuipers et al, ⁴⁵ 1997	UK	38.5/41.8	Outpatient	60 28/32	CBT versus TAU	Individual; 18 sessions fortnightly; 1 hour per session	Hopelessness; self-esteem	BHS, SCQ	–

Continued

Table 1 Continued

Study (year)	Place	Mean age (I/C)	Inpatient or outpatient	Size (n): (I/C)	Study conditions	Form and dose of intervention*	Personal recovery	Instrument	Follow-up† (month)
Lysaker et al, ⁴⁶ 2005	USA	46.7/49.7	Outpatient	50 25/25	CBT versus SS	Group; 8 sessions fortnightly; 40min per session	Hopelessness; self-esteem	BHS, RSE	-
Mortan et al, ⁴⁷ 2011	Manisa	44.0/40.6	Inpatient	12 7/5	CBT versus TAU	Group; 10 sessions fortnightly; 1.5 hours per session	Hopelessness; self-esteem	BHS; RSE	7
Penn et al, ⁴⁸ 2009	USA	41.7/39.6	Outpatient	6 3/3	CBT versus ST	Group; 12 weekly sessions; 1 hour per session	Self-esteem	RSE	9
Premkumar et al, ⁴⁹ 2011	UK	36.1/39.7	Outpatient	43 25/18	CBT versus SC	Group; 19 sessions, weekly/fortnightly; unclear	Self-esteem	RSE	-
Shawyer et al, ⁵⁰ 2012	Australia	40.0/39.6	Outpatient	44 21/23	CBT versus BF	Group; 15 sessions, weekly; 50min per session	Quality of life	Q-LES-Q	3
Steel et al, ⁵¹ 2016	UK	43.8/40.7	Outpatient	31 30/31	CBT versus TAU	Group; 16 sessions over 6 months; unclear	Quality of life	QLS	6
Wykes et al, ⁵² 2005	UK	39.7/39.7	Outpatient	95 45/40	CBT versus TAU	Group; 7 sessions over 10 weeks; unclear	Self-esteem	RSE	4
Tsiachristas et al, ⁵⁶ 2018	UK	40.4/42.9	Outpatient	43 19/24	CBT versus SC	Individual; 8 sessions over 12 weeks; unclear	Quality of life	EQ-5D-5L	3
Wood et al, ⁵⁷ 2018	UK	32.07/35.58	Inpatient	30 15/15	CBT versus PE	Group; 2 sessions over 2 week period; 120min per session	Personal recovery; self-esteem	QPR; SERS	1
Morrison et al, ⁵³ 2018a	UK	23.2/24.4	Outpatient	49 24/25	CBT versus TAU	Group; 26 sessions over 6 months; unclear	Personal recovery; quality of life	QPR; WHO-QoL	12
Morrison et al, ⁵⁴ 2018b	UK	42.8/42.2	Inpatient	475 230/245	CBT versus TAU	Group; 26 sessions over 9 months; 60 min per session	Personal recovery; quality of life	QPR; EQ-5D-5L	8
Pot-Kolder et al, ⁵⁵ 2018	Netherlands	36.5/39.5	Outpatient	116 58/58	CBT versus WL	Individual; 16 sessions over 8-12 weeks; 50min per session	Quality of life	MSAQoL	3

*The information extracted from the primary study were group CBT/individual CBT; number and frequency of CBT sessions; and session length. †The follow-up time was from the endpoint of CBT treatment.

AD, Antipsychotic drugs; BCSS, Brief Core Schema Scales; BF, befriending control group; BHS, Beck Hopelessness Scale; C, control group; CBOS, cognitive-behavioural oriented services; CBT, cognitive-behavioural therapy; CLZ, clozapine; EQ-5D-5L, EuroQol 5 Dimensions 5 Levels; I, intervention group; MSAQoL, Manchester Short Assessment of Quality of Life; MSQoL, Modular System for Quality of Life; NR, Not Report; PE, patient psychoeducation; Q-LES-Q, Quality of Life Enjoyment and Satisfaction Questionnaire; QLS, Quality of Life Scale; QPR, Questionnaire about the Process of Recovery; QoL, quality of life scale; RSE, Rosenberg Self-Esteem Scale; RSQ, Robson Self-Concept Questionnaire; SC, standard care; SCQ, Self Concept Questionnaire; SCS, Social Comparison Scale; SERS, Self-Esteem Rating Scale; SG, support group; SS, standard support; ST, standard treatment; TAU, treatment as usual; TDZ, thioridazine; WEMWBS, Warwick-Edinburgh Mental Well-Being Scale; WL, waiting list group.

Figure 2 Forest plot of the effect of cognitive-behavioural therapy (CBT) on personal recovery

Modular System for Quality of Life (MSQoL) scale and the WHO-QoL psychological well-being subscale. Because of the high heterogeneity ($I^2=91.0\%$, $p<0.001$), it was not appropriate to directly combine the effect sizes; thus, a descriptive analysis was performed. Two studies investigated psychological health with WEMWBS;^{38,39} the pooled effect size was 0.64 (95% CI 0.06 to 1.13; $I^2=30.8\%$; $p<0.01$) and the long-term effect size was 0.38 (95% CI 0.08 to 0.69; $I^2=0.00\%$; $p<0.01$). In a restrictive and high-quality RCT, Bechdolf *et al.*³⁵ investigated the effect of CBT on patients' psychosocial well-being using a subscale of the MSQoL. The results showed that the CBT programme improved the patients' psychological health-related quality of life with a small effect size (Cohen's $d=0.23$). Moreover, the long-term effect of CBT was significant according to data collected 4 months after treatment and it had a moderate effect size (Cohen's $d=0.37$). van der Gaag *et al.*⁴⁰ used the WHO-QoL psychological well-being subscale to reflect the psychological health of patients with schizophrenia. The results showed that CBT significantly improved the patients' QoL, with a large effect size at the end of treatment (Cohen's $d=1.41$).

Shawyer *et al.*⁵⁰ investigated QoL using two subscales from the Quality of Life Enjoyment and Satisfaction Questionnaire: Subjective Feelings and General Activities. In addition, the endpoint effect sizes of the two subscales corresponded to Cohen's d values of 0.02 and 0.43, and the follow-up effect sizes were -0.48 and 0.14 respectively. The study also evaluated the life satisfaction and life enjoyment with special items, and the Cohen's d values were 0.37 and 0.08 with long-term effect sizes of 0.20 and -0.19 respectively. Bechdolf *et al.*³⁵ reported subjective QoL measured with the MSQoL-seven subscale scores at post-treatment and at a 6-month follow-up. The effect sizes of the seven subscales (endpoint, follow-up) were as follows: Physical Health (0.11, 0.21), Vitality (0.01, 0.36), Psychosocial QoL (0.03, 0.37), Affective QoL (0.17, 0.27), Material QoL (0.03, 0.12), Spare Time QoL (0.30, 0.32)

and General QoL (0.03, 0.20). All changes in the effect sizes over time were positive.

Effect size of hope

A total of five studies^{34,36,45-47} investigated the hopelessness levels of patients with schizophrenia using the Beck Hopelessness Scale (BHS), for which higher scores indicate lower hope levels. Egger's test showed that no publication bias existed ($p=0.72$, 95% CI -5.65 to 7.23). The endpoint pooled effect size of the five studies was positive, with an effect size of -1.77 (95% CI -3.29 to -0.25; $I^2=40\%$; $p=0.02$, power=0.89) (figure 3). Regarding the long-term effect of CBT, three studies^{34,36,47} reported follow-up data for over 6 months after the treatment. The results showed that the effect of CBT on improving hope among patients with schizophrenia was uncertain (-0.38, 95% CI -2.78 to 2.02; $I^2=56\%$; $p>0.05$), and the three studies only yielded a power of 0.42. Sensitivity analyses were conducted and after eliminating the studies, no substantial change in the new pooled effect size was observed (figure 4).

Effect size of identity (self-esteem and self-confidence)

A total of eleven studies^{34,39,41,42,45-49,52,57} reported self-esteem. The total sample size of these eleven studies was 584, and the pooled effect size was 1.85 (95% CI 0.69 to 3.01; $I^2=41\%$; $p<0.01$, power=0.98) (figure 5). The result of Egger's test showed that there was no publication bias ($p=0.20$, 95% CI -3.92 to 0.94). For the long-term effect size of CBT, six^{34,39,42,47,48,57} of the ten studies completed a follow-up evaluation and the effect size was -1.21 (95% CI -2.45 to 0.04; $I^2=12\%$; $p>0.05$, power=0.37). Two^{39,57} of those studies reported results 1 month after the end of treatment, which showed an uncertain effect of CBT (3.61, 95% CI -13.89 to 21.11; $I^2=28\%$; $p>0.05$, power=0.11) and the statistical power was 0.11. For the other four studies, the follow-up time was over 6 months. Therefore, we combined the follow-up data from these four studies and it revealed a negative long-term effect,

Figure 3 Forest plot of the effect of cognitive-behavioural therapy (CBT) on quality of life (QoL).

with an effect size of -1.23 (95% CI -2.52 to 0.06 ; $I^2 = 21\%$) to Cohen's d of 0.40 , and the long-term effect size was $p > 0.05$) and a power of 0.49 .

One study³⁹ assessed the patients' self-confidence using the Brief Core Schema Scales (BCSS). The BCSS is designed to assess negative and positive beliefs about oneself and others. The endpoint effect size of positive beliefs corresponded to Cohen's d of 1.14 , but the long-term effect size was only 0.20 at 1 month after treatment. The endpoint effect size of negative beliefs corresponded

Connectedness

The Social Comparison Scale is used to assess the patient's relationship with others, and higher scores indicate a more positive view of oneself in relation to others.³⁹ The effect sizes were 0.79 and 0.33 at the endpoint and follow-up respectively. Klingberg *et al*⁴⁴ investigated the

Figure 4 Forest plot of the effect of cognitive-behavioural therapy (CBT) on hopelessness.

Figure 5 Forest plot of the effect of cognitive-behavioural therapy (CBT) on self-esteem.

effect on social connections of CBT in schizophrenia by the percentage of connections with relatives after CBT treatment, and the result showed significantly more improvement than deterioration regarding social contacts in the CBT group, with 15 of 61 patients exhibiting positive changes after CBT compared with 1 of 60 patients in the control group.

Figure 6 shows the trend of the effect of CBT on personal recovery over time from the end of treatment,

Figure 6 The change in effect size over the follow-up period. Zero on the x-axis indicates the endpoint of treatment.

visually showing the characteristics of CBT long-term effects. Summary of findings tables summarising CoE assessment based on the GRADE approach are shown in table 2.

DISCUSSION

Main findings

There has recently been growing attention surrounding the effects of CBT on personal recovery in patients with schizophrenia. The present review examined the efficacy of CBT across 25 randomised clinical trials that included multiple outcomes of personal recovery over different periods of follow-up. Both the post-treatment and the follow-up effect of CBT on personal recovery measured by QPR were positive and significant, which means that CBT can indeed change the patient's recovery process to some extent. However, due to the limited number of studies, the small sample size and low statistical power, the evidence is not sufficient. We also tested the effect of CBT on the specific components of personal recovery. At post-treatment and during the follow-up period, the effect of CBT on QoL in patients with schizophrenia was uncertain for both pooled groups and subgroups, with p values >0.05 and powers $<80\%$. Thus, a conclusion cannot be drawn as to whether CBT is beneficial for patients' QoL due to the limited number of studies, the small sample sizes and the fair quality of the included controlled trials. Additional sufficient and conclusive evidence is needed in the future.

Table 2 GRADE summary of evidence for the effects of CBT and personal recovery

Certainty assessment							Patients, n		Effect			
Studies, n	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	CBT	TAU	Relative (95% CI)	Absolute (95% CI)	Certainty	Importance
Personal recovery												
3	Randomised trials	Not serious	Not serious	Not serious	Not serious	None	212	230	-	MD 2.27 higher (0.1 higher to 4.45 higher)	⊕⊕⊕⊕ High	Critical
Follow-up (personal recovery)												
3	Randomised trials	Not serious	Not serious	Not serious	Not serious	None	192	219	-	MD 2.62 higher (0.51 higher to 4.47 higher)	⊕⊕⊕⊕ High	Critical
Quality of life												
10	Randomised trials	Serious*	Not serious	Not serious	Serious†	None	456	482	-	MD 0.01 higher (0.12 lower to 0.15 higher)	⊕⊕○○ Low	Critical
Follow-up (quality of life)												
8	Randomised trials	Serious*	Not serious	Not serious	Serious†	None	65	65	-	MD 0.06 higher (0.03 lower to 0.15 higher)	⊕⊕○○ Low	Critical
Hope												
5	Randomised trials	Serious*	Serious§	Serious¶	Serious†	Publication bias strongly suspected**	189	178	-	MD 1.77 lower (3.29 lower to 0.25 lower)	⊕○○○ Very low	Critical
Follow-up (hope)												
3	Randomised trials	Serious*	Serious§	Serious¶	Serious†	None	131	136	-	MD 0.38 lower (2.78 lower to 2.02 lower)	⊕○○○ Very low	Critical
Identity												
10	Randomised trials	Serious*	Serious††	Not serious	Not serious	None	307	277	-	MD 1.85 higher (0.69 higher to 3.01 higher)	⊕⊕○○ Low	Critical
Follow-up (identity)												
6	Randomised trials	Serious*	Not serious	Not serious	Serious†	None	173	174	-	MD 0.11 lower (2.05 lower to 0.04 higher)	⊕⊕○○ Low	Critical

*Allocation concealment bias. Blinding of participants bias and incomplete outcome data bias were observed in some included studies.

†The study included patients who were relatively small and the CIs were wide.

‡Some studies included patients who were relatively small.

§Some included studies reported a positive effect of CBT for hope, whereas other studies failed to find such effect.

¶Some studies use hopeless levels to reflect hope levels.

**Selective reporting.

††Some included studies reported no effect of CBT for identity.

‡‡Some included studies reported no effect of CBT for identity. CBT, cognitive-behavioural therapy; GRADE, Grading of Recommendations Assessment, Development and Evaluation; MD, mean difference; TAU, treatment as usual.

However, for psychological well-being, which is an important component of QoL, the pooled endpoint effect size was significantly large (>0.6), although the long-term effect size decreased to a moderate level. The Subjective Feelings (how much of the time participants feel positive) and General Activities (degree of satisfaction with general activities of life) of patients also showed improvement at post-treatment but decreased at long-term follow-up times. Although this change in effect size with time is in contrary to the trend reported by Bechdolf *et al.*,³⁵ we still believe that CBT is a useful intervention for improving QoL other than psychological health-related QoL in patients with schizophrenia. However, the effect of CBT on QoL in schizophrenia and whether the long-term effect is better or worse than the endpoint should be explored in the future.

The effect of CBT on hope and self-esteem is obvious and positive at post-treatment; however, the long-term effect (over 6 months) is markedly decreased. The self-confidence and relationships of patients with schizophrenia improved little in the CBT group compared with the ST group, and the long-term effect of CBT was the same for hope and self-esteem. These data indicate that the effect of CBT on personal recovery is only sustained for a short time; as time progresses after treatment, the personal recovery of the patients reverts back to the original level. Most studies have only focused on the immediate effects of targeting cognitions about the self and have not tried to determine which specific intervention techniques may change the underlying mechanism.³⁹ We believe that the personal recovery of patients should be a long-term effort, whether in or out of the hospital, and to maximise the effectiveness of CBT, more efforts should be dedicated to continuing CBT intervention at specific time points when its effects are decreased and to determining which factors influence its effects.

None of these studies reported the effect of CBT on meaning and empowerment in patients with schizophrenia. Empowerment includes participating in society in terms of access to employment, education and other valued resources; in terms of interpersonal characteristics, empowerment also means having control over one's life and the recovery, and efforts to achieve greater control and self-efficacy. There are also some scales available for measuring empowerment, especially among patients with psychosis.⁵⁸ Unfortunately, none of the studies included in our research used the empowerment scale as a measure of personal recovery. In the CHIME personal recovery model, meaning is not reported directly, and to the best of our knowledge, there is no specially designed scale for meaning measurement; however, the schizophrenia hope scale designed by Choe⁵⁹ examines positive expectations for the future, confidence in life and the future, and meaning in life, which may help to reflect meaning as an aspect of personal recovery in patients with schizophrenia. Meaning and empowerment can reflect personal recovery to a large extent, and future studies should be designed to verify the effectiveness of

CBT on these components of personal recovery. Moreover, such studies would support the implementation of instruments to measure personal recovery as an outcome.

In the present study, even when strict inclusion criteria were applied to minimise the heterogeneity of the meta-analysis, there was still moderate or even large heterogeneity in some of the outcome analyses. Because of the small number of studies in the high heterogeneity group, the sensitivity analyses and the subgroup analysis were unable to compensate for the heterogeneity; therefore, instead of reporting the results of the meta-analysis, we reported the data as descriptive statistics. Additionally, the outcomes in our study are the best defined main measures for personal recovery despite the moderate heterogeneity of some analyses. The uniformity among estimates of the effect was remarkable given the unavoidable differences in interventions associated with the different individual personnel delivering the behavioural interventions. In addition, we found no statistical evidence of publication bias, therefore such factors are unlikely to have affected our estimates.

Some of the results were not statistically significant, with p values >0.05 . However, the absence of statistical significance should never be interpreted as evidence that an effect is absent. We performed a power analysis to test the reliability of the negative result, and the statistical power was low or very low. According to the significance test ($Z = \frac{M}{SE}$), the effect size (M) is the important factor that determines the p value, and the factors that control the power are the same as those that control the significance⁶⁰ (Chapter 29). Therefore, additional higher power studies with restrictive designs and sufficient sample sizes are needed in the future to confirm the effectiveness of CBT on personal recovery in schizophrenia.

Limitations

There are several limitations of this review. First, some subgroup analyses were not performed due to the limited number of studies. We found that the frequency and number of CBT sessions varied among studies. We initially planned to perform a subgroup analysis according to session design to determine which CBT design corresponds to the best outcome; however, because of the limited number of studies, this subgroup analysis was inappropriate. Therefore, we did not perform this analysis, although this issue could be discussed in future studies aimed at other recovery outcomes. Second, we could not perform a meta-analysis of some outcomes, thus the findings of the effects of CBT are less conclusive and valid. Third, we could not consider all aspects of personal recovery due to the absence of various measures in primary studies. The hope level was represented by the BHS score, which may not be ideal as there is a special hope scale designed for patients with schizophrenia.⁵⁹ Fourth, this review included randomised controlled trials but did not include other study methods/designs or studies that used mixed methods or qualitative exploratory approaches. This limits the comprehensiveness and depth of the understanding of the process,

the perceived benefits and different clinical outcomes of CBT on the personal recovery of patients with schizophrenia. Lastly, the review only included experimental studies published in peer-reviewed journals using English language. This could limit the generalisability and validity of the findings of this review.

Implications

There is insufficient evidence regarding the significant positive long-term effects of CBT on personal recovery outcomes among people with schizophrenia, and more experimental trials with high power are needed in the future. In our study, the QoL and CHIME recovery models were combined to represent personal recovery. However, there are numerous specialised scales designed to measure personal recovery among patients with schizophrenia, and future studies should take these scales into consideration as personal recovery measurement tools.¹³ Other important questions for both research and clinical applications that must be investigated include how long the effect of CBT can be sustained and how to ensure that patients with schizophrenia receive the greatest benefit from CBT intervention in the long term.

CONCLUSION

Our review showed that CBT is a reasonably effective treatment for some aspects of personal recovery among patients with schizophrenia. Our findings revealed improvements in QoL, hope, self-esteem, self-efficacy, and social connections after CBT interventions; however, the effect was relatively immediate and rapidly decreased over time. Therefore, in the future, more studies should focus on the mechanism of CBT for personal recovery and the factors that influence the long-term effects of CBT.

Acknowledgements We thank Lydia Hyman and Kimberly Yasutis, PhD, from American Journal Experts for editing the draft of this manuscript.

Contributors WW and ZY designed the study and contributed substantially to the design of the search strategy. WW and NC searched the literature and extracted the data. DL performed the analysis and interpreted the data. WW wrote the first draft of the manuscript and ZY critically reviewed the manuscript. All authors read and approved the final manuscript.

Funding This work was supported by the National Natural Science Foundation of China (Grant No: 71673070).

Disclaimer The sponsor had no role in the study design, writing of the manuscript, or decision to submit this or future manuscripts for publication.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data generated or analysed during this study are included in this article.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- World Health Organization. *Fact sheet about schizophrenia in media center*. Geneva: World Health Organization, 2018.
- Vos T, Barber RM, Bell B, et al. Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990–2013: a systematic analysis for the global burden of disease study 2013. *Lancet* 2015;386:743–800.
- Saxena S, Funk M, Chisholm D. World health assembly adopts comprehensive mental health action plan 2013–2020. *Lancet* 2013;381:1970–1.
- Health NCCFM. *Schizophrenia: core interventions in the treatment and management of schizophrenia in primary and secondary care (update)*. Social Science Electronic Publishing, 2009.
- National Institute for Health and Care Excellence. *Psychosis and schizophrenia in adults: prevention and management*, 2014. Available: <https://www.nice.org.uk/guidance/cg178>
- Zimmermann G, Favrod J, Trieu VH, et al. The effect of cognitive behavioral treatment on the positive symptoms of schizophrenia spectrum disorders: a meta-analysis. *Schizophr Res* 2005;77:1–9.
- Klingberg S, Wölwer W, Engel C, et al. Negative symptoms of schizophrenia as primary target of cognitive behavioral therapy: results of the randomized clinical studies study. *Schizophr Bull* 2011;37(suppl 2):S98–101.
- YH M. The effects of cognitive behavioral group therapy improving social cognition, self efficacy, relationship function and social skills for chronic schizophrenia. *J Korean Acad Psychiatr Ment Health Nurs* 2017;29.
- Gumley A, Grady M, Murray L, et al. Early intervention for relapse in schizophrenia: results of a 15-month randomized controlled trial of cognitive behavioral therapy. *Psychol Med* 2003;33:419–31.
- Bateman K, Hanrahan L, Turkington D, et al. Cognitive behavioral therapy reduces suicidal ideation in schizophrenia: results from a randomized controlled trial. *Suicide Life Threat Behav* 2007;37:284–90.
- Bellack AS. Scientific and consumer models of recovery in schizophrenia: concordance, contrasts, and implications. *Schizophr Bull* 2006;32:432–42.
- Schmidt-Werner M, Oades L. Recovery: an international perspective. *Epidemiol Psychiatr Soc* 2008;17:128–37.
- Corbin RE, Burger TJ, Vellinga A, et al. The relationship between clinical and personal recovery in patients with schizophrenia spectrum disorders: a systematic review and meta-analysis. *Schizophr Bull* 2017.
- Mead S, Copeland ME. What recovery means to us: consumers' perspectives. *Community Ment Health J* 2000;36:315–28.
- MikeSlade. *Personal recovery and mental illness*. Cambridge: Cambridge University Press, 2009.
- Anthony WA. Recovery from mental illness: the guiding vision of the mental health service system in the 1990s. *J Psychosoc Rehabil Ment Health* 1993;16:11–23.
- Andresen R, Oades L, Caputi P. The experience of recovery from schizophrenia: towards an empirically validated stage model. *Aust N Z J Psychiatry* 2003;37:586–94.
- Leamy M, Bird V, Boutillier CL, et al. Conceptual framework for personal recovery in mental health: systematic review and narrative synthesis. *Br J Psychiatry* 2011;199:445–52.
- Rossi A, Amore M, Galderisi S, et al. The complex relationship between self-reported 'personal recovery' and clinical recovery in schizophrenia. *Schizophr Res* 2017.
- Davidson L. *Living outside mental illness: qualitative studies of recovery in schizophrenia*. New York: New York University Press, 2003.
- Jaeger M, Konrad A, Rueegg S, et al. Patients' subjective perspective on recovery orientation on an acute psychiatric unit. *Nord J Psychiatry* 2015;69:188–95.
- Jerrell JM, Cousins VC, Roberts KM. Psychometrics of the recovery process inventory. *J Behav Health Serv Res* 2006;33:464–73.
- Resnick SG, Rosenheck RA, Lehman AF. An exploratory analysis of correlates of recovery. *PS* 2004;55:540–7.
- Corrigan PW, Salzer M, Ralph RO, et al. Examining the factor structure of the recovery assessment scale. *Schizophr Bull* 2004;30:1035–41.
- World Health Organization. *Promoting recovery in mental health and related services: handbook for personal use and teaching - WHO QualityRights training to act, unite and empower for mental health (pilot version); (WHO/MSD/MHP/17.11)*. Geneva: World Health Organization, 2017.
- Alessandrini M, Lançon C, Fond G, et al. A structural equation modelling approach to explore the determinants of quality of life in schizophrenia. *Schizophr Res* 2016;171:27–34.

- 27 Oles SK, Fukui S, Rand KL, et al. The relationship between hope and patient activation in consumers with schizophrenia: results from longitudinal analyses. *Psychiatry Res* 2015;228:272–6.
- 28 Jahn DR, DeVlyder JE, Drapalski AL, et al. Personal recovery as a protective factor against suicide ideation in individuals with schizophrenia. *J Nerv Ment Dis* 2016;204:827–31.
- 29 Hofmann SG, Asnaani A, Vonk IJJ, et al. The efficacy of cognitive behavioral therapy: a review of meta-analyses. *Cognit Ther Res* 2012;36:427–40.
- 30 Turner DT, van der Gaag M, Karyotaki E, et al. Psychological interventions for psychosis: a meta-analysis of comparative outcome studies. *Am J Psychiatry* 2014;171:523–38.
- 31 Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS ONE* 2009;18:e123.
- 32 Higgins JPT, Green S. Cochrane Handbook for systematic reviews of interventions version 5.1.0. The Cochrane collaboration, 2011. Available: <http://training.cochrane.org/handbook>
- 33 de Paiva Barretto EM, Kayo M, Avrichir BS, et al. A preliminary controlled trial of cognitive behavioral therapy in clozapine-resistant schizophrenia. *J Nerv Ment Dis* 2009;197:865–8.
- 34 Barrowclough C, Haddock G, Lobban F, et al. Group cognitive-behavioural therapy for schizophrenia: randomised controlled trial. *Br J Psychiatry* 2006;189:527–32.
- 35 Bechdolf A, Knost B, Nelson B, et al. Randomized comparison of group cognitive behaviour therapy and group psychoeducation in acute patients with schizophrenia: effects on subjective quality of life. *Aust N Z J Psychiatry* 2010;44:144–50.
- 36 Birchwood M, Michail M, Meaden A, et al. Cognitive behaviour therapy to prevent harmful compliance with command hallucinations (command): a randomised controlled trial. *Lancet Psychiatry* 2014;1:23–33.
- 37 Edwards J, Cocks J, Burnett P, et al. Randomized controlled trial of clozapine and CBT for First-Episode psychosis with enduring positive symptoms: a pilot study. *Schizophr Res Treatment* 2011;2011:1–8.
- 38 Freeman D, Dunn G, Startup H, et al. Effects of cognitive behaviour therapy for worry on persecutory delusions in patients with psychosis (wit): a parallel, single-blind, randomised controlled trial with a mediation analysis. *Lancet Psychiatry* 2015;2:305–13.
- 39 Freeman D, Pugh K, Dunn G, et al. An early phase II randomised controlled trial testing the effect on persecutory delusions of using CBT to reduce negative cognitions about the self and potential benefits of enhancing self confidence. *Schizophr Res* 2014;160:186–92.
- 40 van der Gaag M, Stant AD, Wolters KJK, et al. Cognitive-behavioural therapy for persistent and recurrent psychosis in people with schizophrenia-spectrum disorder: cost-effectiveness analysis. *Br J Psychiatry* 2011;198:59–65.
- 41 Garety PA, Kuipers L, Fowler D, et al. Cognitive behavioural therapy for drug-resistant psychosis. *Br J Med Psychol* 1994;67:259–71.
- 42 Gumley A, Karatzias A, Power K, et al. Early intervention for relapse in schizophrenia: impact of cognitive behavioural therapy on negative beliefs about psychosis and self-esteem. *Br J Clin Psychol* 2006;45:247–60.
- 43 Halperin S, Nathan P, Drummond C, et al. A cognitive-behavioural, group-based intervention for social anxiety in schizophrenia. *Aust N Z J Psychiatry* 2005;34:109–13.
- 44 Klingberg S, Wittorf A, Fischer A, et al. Evaluation of a cognitive behaviourally oriented service for relapse prevention in schizophrenia. *Acta Psychiatr Scand* 2010;121:340–50.
- 45 Kuipers E, Garety P, Fowler D, et al. London-East Anglia randomised controlled trial of cognitive-behavioural therapy for psychosis I: effects of the treatment phase. *Br J Psychiatry* 1997;171:319–27.
- 46 Lysaker PH, Bond G, Davis LW, et al. Enhanced cognitive-behavioral therapy for vocational rehabilitation in schizophrenia: effects on hope and work. *JRRD* 2005;42:673–82.
- 47 Mortan O, Tekinsav Sutcu S, German Kose G. A pilot study on the effectiveness of a group-based cognitive-behavioral therapy program for coping with auditory hallucinations. *Turk Psikiyatri Derg* 2011;22:26–34.
- 48 Penn DL, Meyer PS, Evans E, et al. A randomized controlled trial of group cognitive-behavioral therapy vs. enhanced supportive therapy for auditory hallucinations. *Schizophr Res* 2009;109:52–9.
- 49 Premkumar P, Peters ER, Fannon D, et al. Coping styles predict responsiveness to cognitive behavioural therapy in psychosis. *Psychiatry Res* 2011;187:354–62.
- 50 Shawyer F, Farhall J, Mackinnon R, et al. A randomised controlled trial of acceptance-based cognitive behavioural therapy for command hallucinations in psychotic disorders. *Behav Res Ther* 2012;50:110–21.
- 51 Steel C, Hardy A, Smith B, et al. Cognitive-behaviour therapy for post-traumatic stress in schizophrenia: a randomised controlled trial. *Psychol Med* 2017;47:43–51.
- 52 Wykes T, Hayward P, Thomas N, et al. What are the effects of group cognitive behaviour therapy for voices? a randomised control trial. *Schizophr Res* 2005;77:20–30.
- 53 Morrison AP, Law H, Carter L, et al. Antipsychotic drugs versus cognitive behavioural therapy versus a combination of both in people with psychosis: a randomised controlled pilot and feasibility study. *The Lancet. Psychiatry* 2018;5:411–23.
- 54 Morrison AP, Pyle M, Gumley A, et al. Cognitive behavioural therapy in clozapine-resistant schizophrenia (focus): an assessor-blinded, randomised controlled trial. *The Lancet. Psychiatry* 2018;5:633–43.
- 55 Pot-Kolder H, Geraets CNW, Veling W, et al. Virtual-reality-based cognitive behavioural therapy versus waiting list control for paranoid ideation and social avoidance in patients with psychotic disorders: a single-blind randomised controlled trial. *The Lancet. Psychiatry* 2018;5:217–26.
- 56 Tsiachristas A, Waite F, Freeman D, et al. Cost-Effectiveness of cognitive-behavioural therapy for sleep disorder added to usual care in patients with schizophrenia: the best study. *BJPsych Open* 2018;4:126–35.
- 57 Wood L, Byrne R, Enache G, et al. A brief cognitive therapy intervention for internalised stigma in acute inpatients who experience psychosis: a feasibility randomised controlled trial. *Psychiatry Res* 2018;262:303–10.
- 58 Castelein S, van der Gaag M, Bruggeman R, et al. Measuring Empowerment among people with psychotic disorders: a comparison of three instruments. *PS* 2008;59:1338–42.
- 59 Choe K. Development and preliminary testing of the schizophrenia hope scale, a brief scale to measure hope in people with schizophrenia. *Int J Nurs Stud* 2014;51:927–33.
- 60 Borenstein M, Hedges LV, Higgins JPT, et al. *Introduction to meta-analysis*. Chichester, England: Wiley, 2009.

Weiwei Wang graduated from Xinxiang Medical University of Henan province in 2016. Since 2016, he has been working on successive postgraduate and doctoral programs for Ph.D degree in Harbin Medical University, School of nursing. His research team is currently investigating how to slow down and prevent schizophrenia disability and maintain its social function. In recent years, attention has been paid to the treatment, clinical outcomes and rehabilitation of patients with schizophrenia. They are very interested in the personal recovery of mental illness, the development of clinical assessment tools, and the trajectories of disease development and its relationship with clinical outcomes, brain function and event-related potential studies. His research interest includes mental illness and psychotherapy.

Retraction: Cognitive-behavioural therapy for personal recovery of patients with schizophrenia: a systematic review and meta-analysis

Wang W, Zhou Y, Chai N, *et al*. Cognitive-behavioural therapy for personal recovery of patients with schizophrenia: a systematic review and meta-analysis. *Gen Psychiatr* 2019;32:e100040. doi: 10.1136/gpsych-2018-100040.

This paper has been retracted after errors were found in the original data of the study that affect the main result of the paper. The authors are going to check and renew all the processes of the screening and data collection. Therefore, we are in agreement with the author that this paper should be retracted pending further analysis

OPEN ACCESS

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

General Psychiatry 2019;32:e100040ret. doi:10.1136/gpsych-2018-100040ret